

Occurrence of *Campylobacter jejuni* and *C. coli* on broiler carcasses after chilling in southern Brazil¹

Gustavo Perdoncini², Yuli M. Sierra-Arguello², Leonardo M. Lima², Michele M. Trindade², Marcos José Pereira Gomes², Luciana R. dos Santos³, Verônica Schmidt⁴ and Vladimir Pinheiro do Nascimento²

ABSTRACT.- Perdoncini G., Sierra-Arguello Y.M., Lima L.M., Trindade M.M., Gomes M.J.P., Santos L.R., Schmidt V. & Nascimento V.P. 2015. **Occurrence of *Campylobacter jejuni* and *C. coli* on broiler carcasses after chilling in southern Brazil.** *Pesquisa Veterinária Brasileira* 35(4):349-352. Centro de Diagnóstico e Pesquisa em Patologia Aviária, Faculdade de Veterinária, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 8824, Porto Alegre, RS 91540-000, Brazil. E-mail: gustavo.perdoncini@yahoo.com

Campylobacter jejuni and *C. coli* have been associated with gastrointestinal disorders in human beings, due mainly to the consumption of chicken meat. Despite control measures for reducing contamination by these bacteria, the detection of *Campylobacter* in carcasses after chilling remains high. A total of 105 carcasses were assessed by the horizontal detection method in five federally inspected slaughterhouses in southern Brazil in 2012 and in the first three months of 2013. *Campylobacter* was isolated in 37.1% of the carcasses, of which 97.5% contained *C. jejuni* and 2.5% were infected by *C. coli*. The rate of positive carcasses across the slaughterhouses ranged from 0 to 71.4%. Determining the occurrence of *Campylobacter* among flocks is crucial for estimating the microbial load at specific points along the slaughtering process and for minimizing the risk of contamination of end products by *Campylobacter*.

INDEX TERMS: Campylobacteriosis, cross-contamination, broilers, zoonosis.

RESUMO.- [*Campylobacter jejuni* e *C. coli* em carcaças de frangos após a refrigeração por imersão no sul do Brasil.] *Campylobacter jejuni* e *C. coli* têm sido associados a problemas gastroentéricos em seres humanos principalmente devido ao consumo de carne de frango. Embora medidas de controle sejam realizadas para reduzir a contaminação por estas bactérias, a identificação de *Campylobacter* em carcaças após a refrigeração por imersão é alta. Foram analisadas 105 carcaças pelo método de detecção horizontal em cinco abatedouros sob Inspeção Federal no sul do Brasil em 2012 e nos três primeiros meses de 2013. *Campylobacter* foi isolada em 37,1% das carcaças analisadas,

as quais 97,5% foram identificados como *C. jejuni* e 2,5% como *C. coli*. A ocorrência de carcaças positivas entre matadouros variou de zero a 71,4%. O conhecimento sobre a ocorrência de *Campylobacter* entre os lotes é fundamental para estimar a extensão da carga microbiana em pontos específicos do abate e consequentemente minimizar o risco de contaminação por *Campylobacter* em produtos finais de frangos.

TERMOS DE INDEXAÇÃO: Campilobacteriose, contaminação cruzada, frangos, zoonose.

INTRODUCTION

Thermotolerant *Campylobacter* spp. are the causative agents of campylobacteriosis (Rosenquist et al. 2003, Butzler 2004), a zoonotic disease (WHO 2000) transmitted to human beings mainly by the consumption of contaminated poultry products (Dickins et al. 2002). In the European Union (EFSA 2012) and in the United States (CDC 2013) *Campylobacter jejuni* and *Campylobacter coli* are often associated with gastroenteritis, and *C. jejuni* is also implicated in Guillain-Barré syndrome (Vucic et al. 2009). The rise

¹ Received on September 16, 2014.

Accepted for publication on April 9, 2015.

² Centro de Diagnóstico e Pesquisa em Patologia Aviária (CDPA), Faculdade de Veterinária, Universidade Federal do Rio Grande do Sul (UFRGS), Av. Bento Gonçalves 8824, Porto Alegre, RS 91540-000, Brazil. *Corresponding author: gustavo.perdoncini@yahoo.com

³ Faculdade de Veterinária e Agronomia, Universidade de Passo Fundo (UPF), BR-285 Km 171, Cx. Postal 611, Passo Fundo, RS 99001-970, Brazil.

⁴ Faculdade de Veterinária, UFRGS, Av. Bento Gonçalves 9090, Porto Alegre, RS 91540-000.

in the incidence of campylobacteriosis in several countries, especially after 1990, has caused a lot of concern due to the growing morbidity and mortality rates among children and immunocompromised individuals (WHO 2000). This has encouraged research studies in an attempt to minimize the number of positive flocks and to have a better understanding of the intrinsic characteristics of these bacteria.

Seeking to reduce the impact of this pathogen on public health, research studies have investigated the factors that contribute to contamination and persistence of *Campylobacter* spp. on poultry farms. The direct contact of birds with feces, untreated water, rodents, wild birds, or vectors such as *Alphitobius diaperinus* (darkling beetles) infected with this bacterium may contaminate the flocks to be slaughtered (Eberle et al. 2013).

At meat-packing plants, contaminated feces and feathers help disseminate *Campylobacter* in the slaughtering process. Defeathering is critical for contamination as the concentration of *Campylobacter* increases at this stage, as well as in evisceration (Herman et al. 2003, Rosenquist et al. 2006, Figueroa et al. 2009), and measures such as rinsing and chilling of carcasses are alternatives for reducing the incidence of *Campylobacter* (Reich et al. 2008, Figueroa et al. 2009).

However, contamination may occur at any time during the slaughtering process, for instance, in scalding, defeathering, and chilling (Rosenquist et al. 2006, Allen et al. 2007, Figueroa et al. 2009). A study that investigated previous colonization of carcasses by *Campylobacter* spp. detected contamination with *C. jejuni* and *C. coli* in broiler carcasses in two out of five previously negative flocks (Allen et al. 2007).

Given the importance of this bacterium to public health and to poultry production, this paper assesses the occurrence of *C. jejuni* and *C. coli* in broiler carcasses after chilling at meat-packing plants in southern Brazil.

MATERIALS AND METHODS

A cross-sectional observational study was conducted and convenience samples were used (Thrusfield 2004). In total, 105 broiler carcasses were collected after chilling from five federally inspected slaughterhouses in southern Brazil in 2012 and in the first three months of 2013. The samples were stored in isothermal ice boxes and taken to the laboratory for identification of *C. jejuni* and *C. coli* by the horizontal method (ISO 10272-1:2006). The chilled carcasses were rinsed in sterile polyethylene bags containing 400 mL with buffered peptone water (Oxoid®) and aliquots of 1 mL were obtained and homogenized in 9 mL of Bolton broth - Oxoid® (1:9), which were then incubated in a microaerophilic environment (5% O₂, 10% CO₂ and 85% N₂) at 41.5±0.5°C for 48 hours. Thereafter, 100 µL of the broth suspension was filtered in acetate membrane measuring 0.65 µm x 47 mm (Skirrow 1977, Bolton 1982) plated onto mCCD agar (CM739, Oxoid®) for 30 minutes, followed by incubation under the conditions described earlier. *Campylobacter*-compatible colonies were grown on 7% sheep blood agar.

For positive control of isolation and PCR assays were used two references strains, *C. jejuni* subsp. *jejuni* ATCC 29428 and *C. coli* ATCC 43478. The negative control for PCR assay was used *Arco-bacter* sp. strain.

The DNA of the colonies was obtained by thermal extraction

and used for multiplex PCR as proposed by Denis et al. (1999) with some changes, which allows distinguishing *C. jejuni* from *C. coli*. Three primer pairs were used for each reaction, based on the sequence of primers for the common 16S rRNA region between the species: MD16S1, (16S rRNA) *F*^a - ATCTAATGGCTTAACCATTAAC e *R*^b - GGACGGTAACTAGTTTAGTATT with 857 pb. The identification of *C. jejuni* and *C. coli* was based on the genes *mapA* *F*^a - CTATTTTATTTTGAGTGCTTG and *R*^b - GCTTTATTTGCCATTGTTT-TATTA with 589 pb and *ceuE* *F*^a - AATTGAAAATTGCTCCAATATG and *R*^b - TGATTTTATTTTGTAGCAGCG with 462 pb, respectively.

The amplification conditions were as follows: denaturation (10 minutes at 95°C), 35 cycles of denaturation (30 seconds at 95°C), annealing (1 minute and 30 seconds at 59°C), extension (1 min at 72°C), followed by final extension (10 minutes at 72°C). All reactions were performed in a thermocycler (Biocycler - Peltier Thermal Cycler MJ96+MJ96G) and visualized by 1.5% agarose gel electrophoresis stained with ethidium bromide and by ultraviolet transillumination imaging.

RESULTS AND DISCUSSION

Campylobacter was detected in 37.1% of the 105 analyzed carcasses, showing prevalence of *C. jejuni* (97.5%) followed by *C. coli* (2.5%). Both species were detected in two samples (5.1%).

These rates are lower than those reported for European Union countries in 2008 (EFSA 2010), where 75.3% of chilled carcasses were positive for *Campylobacter* spp., with a range from 4.9% (Estonia) to 100% (Luxembourg) and detection of *C. jejuni* in 60.8% of the flocks, followed by *C. coli* and *C. lari* with 41.5%. Carcasses before final rinsing may have a high incidence (up to 100%) of *Campylobacter*, but contamination at this stage is significantly reduced by rinsings and later refrigeration (Son et al. 2007).

In this study the rates of *Campylobacter*-positive samples varied from 0 to 71.4% in chilled carcasses at the five meat-packing plants. In fact, the bacterium was not isolated from only one of the plants (Table 1). These differences can be attributed to the procedures carried out at meat-packing plants, their characteristics, quality management, and minimum operating health standards throughout the slaughtering process, indicating that any problems with these items will be closely related to the final contamination of carcasses (Habib et al. 2012). Another important factor concerns positive flocks, which influence the contamination of equipment and, consequently, the microbiological quality of the end product (Smith et al. 2007, Malher et al. 2011). Consonant with these findings, Kuana et al. (2008) assessed 22 broiler flocks and observed that only

Table 1. Occurrence of *Campylobacter jejuni* and *C. coli* at meat-packing plants and in carcasses after chilling

Meat-packing plant	Positive carcasses	(%)	Negative carcasses	(%)	Total
A	5*	27,78	13	72,22	18
B	12**	50	12	50	24
C	12	57.14	9	42.86	21
D	10	71.42	14	28.58	24
E	0	0	18	100	18
TOTAL	39(37.14%)		66(62.86%)		105

*One positive sample only for *Campylobacter coli*; **Two positive samples for both species. The other isolates were identified as *C. jejuni*.

four of them were not positive for *Campylobacter* spp. before slaughter but that changed during the processing stage with the isolation of the bacterium in chilled carcasses.

The increase in carcass contamination is often related to fowl size and to equipment setup. It should be noted, though, that even when the equipment is properly regulated, there may be intestinal rupture in broilers and contamination of the slaughter line. Two different approaches can be used for reducing and/or eliminating the bacterium: adoption of specific control measures for the reduction of *Campylobacter*-positive farms at the primary level, and actions for minimizing the contamination of ready-to-sell carcasses at the industry level (Hald et al. 2000).

Rinsing and refrigeration of carcasses are alternatives for controlling contamination during the slaughtering process, while the immersion of carcasses in water at 75°C for 30 seconds (Corry et al. 2007) or the use of vapor at 90°C for 12 seconds (Whyte et al. 2003) after chilling yielded *Campylobacter* reduction rates of 1.6 log₁₀ cfu/cm² and 1.3 log₁₀ cfu/g, respectively. The use of organic acids in the chilling process has been assessed and has reduced contamination, but these products have an impact on the sensory features of carcasses (Nagel et al. 2013).

Investment in logistics and verification of flocks for the presence of *Campylobacter* before slaughter help reduce cross-contamination (Nauta & Havelaar 2008) and would be an alternative for avoiding putting negative and positive flocks together. Nevertheless, this strategy is not successful when traditional detection methods are used, as such methods require at least 48 hours for confirmation of positive results. Once negative flocks are identified in the field, it is necessary to ensure that broilers will not be contaminated during transportation from the farms to the slaughterhouse (Hansson et al. 2005). However, a study conducted in the Netherlands by Nauta et al (2009) assessed 62 broiler flocks and did not find significant support for the implementation of control strategies targeted at the identification of positive flocks in the field and at slaughtering time. The correlation between the contamination of feces and of breast cuts by *Campylobacter* suggests these assessed criteria are not good indicative signs of human exposure to *Campylobacter*.

Only the identification of the bacterium in the field would not be an indication of possible human campylobacteriosis, due to the influence of slaughtering technology, which makes the detection of *Campylobacter* in carcasses an essential tool for evaluating the risk of contamination. The adoption of measures that can reduce the number of positive carcasses goes beyond slaughtering logistics (Evers 2004), as other measures such as improvements in scalding, in the configuration of defeathering equipment and techniques, in evisceration, and additional rinsings of carcasses are also important.

The inclusion of qualitative microbiological analysis of carcasses is paramount in checking for the presence of pathogens after the processing stage. This allows estimating the microbial load at specific points along the process, thereby minimizing the incidence of *Campylobacter* in carcasses after chilling and improving the microbiological quality of chicken meat.

Acknowledgments. To CNPq (Brazilian National Council for Scientific and Technological Development) and CAPES (Coordination for the Improvement of Higher Education Personnel) for the financial support and fellowship grants. We also thank to professor Marcos José Pereira Gomes, head of the Laboratory of Veterinary Bacteriology of Universidade Federal do Rio Grande do Sul, and to the CDPA (Diagnostic and Research Center for Avian Pathology) for their technical and structural support.

REFERENCES

- Allen M.V., Bull S.A., Corry J.E.L., Domingue G., Jørgensen F., Frost J.A., Whyte R., Gonzalez A., Elviss N. & Humphrey T.J. 2007. *Campylobacter* spp. contamination of chicken carcasses during processing in relation to flock colonisation. *Int. J. Food Microbiol.* 113:54-61.
- Butzler J.P. 2004. *Campylobacter*, from obscurity to celebrity. *Clin. Microbiol. Infect.* 10:868-876.
- Bolton F.J. & Robertson L. 1982. A selective medium for isolating *Campylobacter jejuni/coli*. *J. Clin. Pathol.* 35:462-467.
- CDC 2013. Incidence and Trends of Infection with Pathogens Transmitted Commonly Through Food - Foodborne Diseases Active Surveillance Network, 10 U.S. Sites, 1996-2012. *Morbidity and Mortality Weekly Report.* 19:283-287.
- Corry J.E.L., James S.J., Purnell G., Pinto C.S., Chochois Y., Howell M. & James C. 2007. Surface pasteurisation of chicken carcasses using hot water. *J. Food Eng.* 76:913-919.
- Dickins M.A., Franklin S., Stefanova R., Schutze G.E., Eisenach K.D., Wesley I. & Cave M.D. 2002. Diversity of *Campylobacter* isolates from retail poultry carcasses and from humans demonstrated by pulse-field gel electrophoresis. *J. Food Protect.* 65:957-962.
- Denis M., Soumet C., Rivoal K., Ermet G., Blivet D., Salvat G. & Colin P. 1999. Development of a m-PCR assay for simultaneous identification of *Campylobacter jejuni* and *C. coli*. *Lett. Appl. Microbiol.* 29:406-410.
- Eberle K.N., Davis J.D., Purswell J.P., Parker H.M., McDaniel C.D. & Kiess A.S. 2013. A One Year Study of Newly Constructed Broiler Houses for the Prevalence of *Campylobacter*. *Int. J. Poultry Sci.* 12:29-36.
- EFSA 2010. Analysis of the baseline survey on the prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on broiler carcasses in the EU, 2008. Part A: *Campylobacter* and *Salmonella* prevalence estimates. *Eur. Food Safety Authority J.* 8:1-99.
- EFSA 2012. Scientific Opinion on a review on the European Union Summary Reports on trends and sources zoonoses, zoonotic agents and food borne outbreaks in 2009 and 2010: specifically for the data on *Salmonella*, *Campylobacter*, verotoxigenic, *Escherichia coli*, *Listeria monocytogenes* and foodborne outbreaks. *Eur. Food Safety Authority J.* 10:1-25.
- Evers E.G. 2004. Predicted quantitative effect of logistic slaughter on microbial prevalence. *Prev. Vet. Med.* 65:31-46.
- Figueroa G., Troncoso M., López C., Rivas P. & Toro M. 2009. Occurrence and enumeration of *Campylobacter* spp. during the processing of Chilean broilers. *BMC Microbiol.* 9:1-6.
- Habib I., Berkvens D., Zutter L. De, Dierick K., Huffel X.V., Speybroeck N., Geeraerd A.H. & Uyttendaele M. 2012. *Campylobacter* contamination in broiler carcasses and correlation with slaughterhouses operational hygiene inspection. *Food Microbiol.* 29:105-112.
- Hald B., Wedderkopp A. & Madsen M. 2000. Thermophilic *Campylobacter* spp. in Danish broiler production: A cross-sectional survey and a retrospective analysis of risk factors for occurrence in broiler flocks. *Avian Pathol.* 29:123-131.
- Hansson I., Ederoth M., Andersson L., Vagsholm I. & Olsson Engvall E. 2005. Transmission of *Campylobacter* spp. to chickens during transport to slaughter. *J. Appl. Microbiol.* 99:1149-1157.
- Herman L., Heyndrickx M., Grijspeerdt K., Vandekerckhove D., Rollier I. & De Zutter L. 2003. Routes for *Campylobacter* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiol. Infect.* 131:1169-1180.
- ISO 2006. International Standard Organization 10272-1:2006 describes a horizontal method for the detection of *Campylobacter* spp.

- Kuana S.L., Santos L.R., Rodrigues L.B., Salle C.T.P.S., Moraes H.L.S. & Nascimento V.P. 2008. Occurrence and Characterization of *Campylobacter* in the Brazilian Production and Processing of Broilers. *Avian Dis.* 88:680-684.
- Malher X., Simon M., Charnay V., Déserts R.D., Lehébel A. & Belloc C. 2011. Factors associated with carcass contamination by *Campylobacter* at slaughterhouse in cecal-carrier broilers. *Int. J. Food Microbiol.* 150:8-13.
- Nagel G.M., Bauermeister L.J., Bratcher C.L., Singh M. & McKee R.S. 2013. *Salmonella* and *Campylobacter* reduction and quality characteristics of poultry carcasses treated with various antimicrobials in a post-chill immersion tank. *Int. J. Food Microbiol.* 165:281-286.
- Nauta M.J. & Havelaar A.H. 2008. Risk-based standards for *Campylobacter* in the broiler meat chain. *Food Control* 19:372-381.
- Nauta M.J., Van der Wal F.J., Putirulan F.F., Post J., Van der Kassteele J. & Bolder N.M. 2009. Evaluation of the "testing and scheduling" strategy for control of *Campylobacter* in broiler meat in The Netherlands. *Int. J. Food Microbiol.* 134:216-222.
- Reich F., Atanassova V. & Klein G. 2008. The effects of *Campylobacter* numbers in caeca on the contamination of broiler carcasses with *Campylobacter*. *Int. J. Food Microbiol.* 127:116-120.
- Rosenquist H., Nielsen N.L., Sommer H.M., Nørrung B. & Christense B.B. 2003. Quantitative risk assessment of human campylobacteriosis associated with thermophilic *Campylobacter* species in chickens. *Int. J. Food Microbiol.* 83:87-103.
- Rosenquist H., Sommer H.M., Nielsen N.L. & Christensen B.B. 2006. The effect of slaughter operations on the contamination of chicken carcasses with thermotolerant *Campylobacter*. *Int. J. Food Microbiol.* 108:226-232.
- Skirrow M.B. 1977. *Campylobacter enteritis*: a "new" disease. *Brit. Med. J.* 2:9-11.
- Smith D.P., Northcutt J.K., Cason J.A., Hinton Jr A., Buhr R.J. & Ingram K.D. 2007. Effect of external or internal fecal contamination on numbers of bacteria on pre-chilled broiler carcasses. *Poultry Sci.* 86:1241-1244.
- Son I., Englen M.D., Berrang E.B., Fedorka-Cray P.J. & Harrison M.A. 2007. Prevalence of *Arcobacter* and *Campylobacter* on broiler carcasses during processing. *Int. J. Food Microbiol.* 113:16-22.
- Thrusfield M. 2004. *Epidemiologia Veterinária*. Roca, São Paulo. 556p.
- Vucic S., Kierman M. & Comblath D.R., 2009. Guillain-Barré syndrome: An update. *J. Clin. Neurosci.* 16:733-741.
- Whyte P., McGill K. & Collins J.D. 2003. An assessment of steam pasteurisation and hot water immersion treatments for the microbiological decontamination of broiler carcasses. *Food Microbiol.* 20:111-117
- World Health Organization - WHO. *Campylobacter*. 2000. Disponível em <<http://www.who.int/mediacentre/factsheets/fs255/en/print.html>> Access in Aug. 2013.