

Production and characterization of monoclonal antibodies against *Campylobacter fetus* subsp. *venerealis*¹

Telma M. Alves², Luiz G.D. Heneine³, Bárbara S. Araújo², Luciana M. Silva⁴, Patrícia C. Campos³, Márcia S. Hermogenes⁴ and Andrey P. Lage^{2*}

ABSTRACT- Alves T.M., Heneine L.G.D., Araújo B.S., Silva L.M., Campos P.C., Hermogenes M.S. & Lage A.P. 2012 **Production and characterization of monoclonal antibodies against *Campylobacter fetus* subsp. *venerealis*.** *Pesquisa Veterinária Brasileira* 32(7)640-644. Laboratório de Bacteriologia Aplicada, Departamento de Medicina Veterinária Preventiva, Escola de Veterinária, Universidade Federal de Minas Gerais. Av. Antônio Carlos 6627, Cx. Postal 567, Belo Horizonte, MG 30123-970, Brazil. E-mail: alage@vet.ufmg.br.

Myeloma cells Sp2/0-Ag14 and spleen cells from BALB/c mouse immunized with sonicated *Campylobacter fetus* subsp. *venerealis* NCTC 10354 were fused with polyethylene glycol (PEG) for the selection of clones producing antibodies. Clones were obtained by limiting dilution and screened for the production of specific antibodies to *C. fetus* subsp. *venerealis* NCTC 10354 by indirect ELISA and western blot against a panel of bacteria: *C. fetus* subsp. *venerealis* NCTC 10354, *C. fetus* subsp. *fetus* ADRI 1812, *C. sputorum* biovar *sputorum* LMG 6647, *C. lari* NCTC 11352, and *Arcobacter skirrowii* LMG 6621 for the ELISA and *C. fetus* subsp. *venerealis* NCTC 10354 and *C. sputorum* biovar *sputorum* LMG 6647 for the western blotting. Fifteen clones producing monoclonal antibodies (MAbs) anti-*C. fetus* subsp. *venerealis* of the IgM (1) and IgG (14) classes were further screened for species-specificity. Four clones of the 15 obtained were producers of species-specific monoclonal antibodies (MAbs): two were specific for *C. fetus* subsp. *venerealis* and two were specific for *C. fetus* subsp. *fetus*. None of the clones were reactive against *C. sputorum* biovar *sputorum* LMG 6647. All clones recognized a protein with molecular mass of approximately 148 kDa from lysed *C. fetus* subsp. *venerealis* NCTC 10354.

INDEX TERMS: Monoclonal antibodies, *Campylobacter fetus* subsp. *venerealis*, cattle, bovine genital campylobacteriosis.

RESUMO.- [Produção e caracterização de anticorpos mono-específicos contra *Campylobacter fetus* subsp. *venerealis*.] Para a produção de anticorpos monoclonais contra *Campylobacter fetus* subsp. *venerealis* foram utilizadas as linhagens de células de mieloma Sp2/0-Ag14 e células de baço de camundongos BALB/c imunizados com soni-

cado de *C. fetus* subsp. *venerealis* NCTC 10354. A detecção dos anticorpos monoclonais foi realizada por ELISA indireto utilizando antígeno sonicado de *C. fetus* subsp. *venerealis* NCTC 10354. A clonagem foi realizada por diluição limitante e os clones foram caracterizados por ELISA indireto utilizando um painel de bactérias escolhidas em função da prevalência e habitats: *C. fetus* subsp. *venerealis* NCTC 10354, *C. fetus* subsp. *fetus* ADRI 1812, *C. sputorum* biovar *sputorum* LMG 6647, *C. lari* NCTC 11352 e *Arcobacter skirrowii* LMG 6621; e no "western blotting" utilizando antígenos sonicados de *C. fetus* subsp. *venerealis* NCTC 10354 e *C. sputorum* biovar *sputorum* LMG 6647. Foram obtidos 15 clones produtores de anticorpos anti- *C. fetus* subsp. *venerealis* das classes IgM (1) e IgG (14). Quatro clones dentre os 15 clones obtidos foram produtores de anticorpos monoclonais espécie-específicos: dois clones reagiram com maior especificidade contra *C. fetus* subsp. *venerealis* NCTC 10354 e dois clones reagiram com maior especificidade

¹ Received on December 13, 2011.

Accepted for publication on March 10, 2012.

² Laboratório de Bacteriologia Aplicada, Núcleo de Pesquisa em Saúde Animal, Departamento de Medicina Veterinária Preventiva, Escola de Veterinária, Universidade Federal de Minas Gerais (UFMG), Av. Antônio Carlos 6627, Cx. Postal 567, Belo Horizonte, MG 30123-970, Brazil. *Corresponding author: alage@vet.ufmg.br

³ Laboratório de Imunologia e Bioprodutos, Fundação Ezequiel Dias (Funed), Rua Conde Pereira Carneiro 80, Bairro Gameleira, Belo Horizonte, MG 30510-010.

⁴ Laboratório de Biologia Celular e Molecular, Funed, Belo Horizonte, MG.

contra *C. fetus* subsp. *fetus* ADRI 1812. Nenhum dos clones reagiu contra *C. sputorum* biovar *sputorum* LMG 6647, comprovando a especificidade dos anticorpos monoclonais testados. Todos os clones reconheceram uma proteína de massa molecular de aproximadamente 148 kDa no sonicação de *C. fetus* subsp. *venerealis* NCTC 10354.

TERMOS DE INDEXAÇÃO: Anticorpos monoclonais, *Campylobacter fetus* subsp. *venerealis*, bovinos, campilobacteriose genital bovina.

INTRODUCTION

Campylobacter fetus is a Gram-negative, microaerophilic bacterium, classified into two closely related subspecies, *C. fetus* subsp. *fetus* and *C. fetus* subsp. *venerealis*. It is the causative agent of different diseases in man, cattle and sheep. Genital campylobacteriosis is a sexually transmitted disease, with a worldwide distribution, that affects cattle resulting in great economic loss due to the reproductive problems it produces (Blaser et al. 2008, Alves et al. 2011).

The habitat of *C. fetus* subsp. *venerealis* is the genital tract of cattle where the microorganism is found predominantly in the epithelial crypts of prepuce and penis and the epithelia of vagina and uterus, and is responsible for repeated breedings and abortions. In contrast, the habitat of *C. fetus* subsp. *fetus* is the gastrointestinal tract, and is the cause of abortion in sheep and sporadic abortion in cattle (Stoessel 1982, Blaser et al. 2008).

In Brazil, about 20% of beef bulls are infected with *C. fetus* and 50.8% of the beef herds have infected animals, but in some regions those prevalences were higher than 50% and 85%, respectively, indicating a wide distribution of *C. fetus* subsp. *venerealis* in the country (Alves et al. 2011).

In 1975, in England, Köhler and Milstein reported the production of monoclonal antibodies (MAbs) with high specificity for a given epitope. From then on, antibodies with homogeneous and high specificity for a single epitope could be obtained. The aim of the present study was to produce and characterize MAbs against *C. fetus* subsp. *venerealis*.

MATERIAS AND METHODS

Animals. Female BALB/c mice aged between 4 and 6 weeks were purchased from the Centro de Bioterismo do Instituto de Ciências Biológicas (ICB) from Universidade Federal de Minas Gerais (UFMG) and used for the MAbs production. Specific pathogen free (SPF) female Swiss mice, used for the removal of peritoneal macrophages, were obtained from the animal facility of Fundação Ezequiel Dias (Funed).

The project was approved by the Ethics Committee in Animal Experimentation (Cetea/UFMG) by the protocol nº 124/04. All animal manipulation followed humane rules for animal handling (AVMA 2001).

Bacterial strains and culturing conditions. *Arcobacter skirrowii* LMG6621 (Laboratorium voor Microbiologic, Rijksuniversiteit Gent, Belgium), *Campylobacter fetus* subsp. *fetus* ADRI 1812 (Animal Diseases Research Institute, Canada), *C. fetus* subsp. *venerealis* NCTC10354 (National Type Culture Collection, England), *C. hyointestinalis* subsp. *hyointestinalis* LCDC17398 (Laboratory Center for Disease Control, Canada), *C. lari* NCTC 11351, and *C. sputorum* biovar *sputorum* LMG 6647 were cultured in Brain He-

art Infusion Agar (BHI, Difco, EUA), supplemented with 10% horse blood, under microaerophilic condition (5% O₂, 5% H₂, 10% CO₂ and 80% N₂) at 37°C for 48h (Debruyne et al. 2008). Bacterial cells were harvested and suspended in phosphate buffered saline (PBS) pH 7.4, standardized to the McFarland nephelometer 10 scale and sonicated (Sonics Vibra Cell™, USA) in ice for 6 cycles of 30 second each, with 1 minute intervals. The frequency used was 50 kHz and a bacterial lysis of 60% was established as optimal. The sonicated antigen was stored at -80°C. The concentration of proteins was determined by the method of Lowry et al. (1951).

Immunization of mice. BALB/c female mice were immunized with the sonicated antigen of the *C. fetus* subsp. *venerealis* strain NCTC 10354, emulsified in Freund's complete adjuvant (Sigma, USA) (first inoculation) and Freund's incomplete (Sigma, USA) (second and third inoculations) in the proportion of 1:2. The inoculated volumes were of 500 µL containing 50µg of antigen and the intraperitoneal route was employed at days 0, 15 and 30. A fourth inoculation was performed by the intravenous or subcutaneous route, with 50µg of the antigen in 100µL of PBS without adjuvant, four days before removal of the spleen, for B cells synchronization (Harlow & Lane 1988).

Preparing spleen cells. BALB/c mice were sacrificed by cervical dislocation (AVMA 2001) four days after the final boost, and spleen was removed and washed three times with RPMI 1640 medium (Invitrogen, USA) supplemented with 10mg/mL tetracycline (Sigma, USA), 1mg/mL amphotericin B (Bristol-Myers Squibb, Brazil) and 40mg/mL gentamicin sulfate (Sigma, USA). The cells were expelled from the spleen by maceration and 5 mL of lysis buffer (ammonium chloride 0.16 M, Tris 0.17 M) were added to the cells, which stand for 5 minutes at room temperature (Hurrell 1988). The cells were then transferred to a tube and centrifuged once for 5 minutes, at 200 x g, at 25°C. The pellet was resuspended and part of the cells was frozen according to Marusich (1988) with modifications: dimethyl sulfoxide (DMSO, Sigma, USA), RPMI 1640 and fetal bovine serum (FBS, Gibco, USA) were added to the cells prior to storage at -80°C, at a cell concentration of 1.0 x 10⁷ cells/mL.

Culture of macrophages from Swiss SPF mice. Swiss SPF mice were sacrificed by cervical dislocation (AVMA 2001) and the peritoneal cavity was rinsed three times with warm RPMI 1640 supplemented with HAT (100mL RPMI 1640 medium + 2.4mL HAT - Sigma, USA - plus 15% FBS, 16.1g/mL bovine insulin - Sigma, USA, 10 mg/mL tetracycline, 1mg/mL amphotericin B and 40mg/mL gentamicin sulfate).

Culture of myeloma line Sp2/0-Ag14. The Sp2/0-Ag14 myeloma cell line (ATCC CRL 1581, American Type Culture Collection, USA, Köhler & Milstein 1975) was previously thawed, cultured in pre-fusion medium (100mL RPMI-1640 supplemented with 10% FBS and 20µg/mL 8-azaguanine, Sigma, USA), in an incubator at 37°C with 5% CO₂ for seven days. Then, the cells were cultured for seven more days in basal medium (RPMI 1640 supplemented with 10% FBS).

Fusion. Fusion was performed according to Harlow & Lane (1988) using fresh and thawed spleen cells (Marusich 1988). Fourteen days after fusion, the screening of hybridomas was initiated using indirect ELISA to detect antibodies against *C. fetus* subsp. *venerealis* (Harlow & Lane 1988, Coligan et al. 1998).

Indirect ELISA to detect *Campylobacter fetus* subsp. *venerealis*. Microtiter plates with flat bottom (PoliSorp™, Nunc, Denmark) were coated with the sonicated antigen of *C. fetus* subsp. *venerealis* NCTC 10354, at 3ng/µL, diluted in carbonate/bicarbonate buffer 0.05 M pH 9.6, at 4°C, overnight. The volume for all assay steps was of 100µL/well. After washing the plates with washing solution (saline-NaCl + 0.05% Tween 20) they were blocked with blocking buffer - 0.1M PBS pH 7.4 + 3% bovine serum albumin and 0.05% Tween 20, at 37°C, for 1 h.

The hybridoma supernatant were added to the plates, and incubated at 37°C for 90 minutes. Plates were washed as previous described and conjugate (anti-mouse IgG conjugated with peroxidase, Sigma, USA), diluted in 1:20,000 in incubation buffer (0.1M PBS + 0.1% Tween 20), was added and the plates incubated for 1 h at 37°C.

The substrate ortophenilenediamine (OPD, Sigma, USA) was added at 40mg/100mL, in 0.15 M citrate buffer pH 5.0, and the plates incubated for 1 h at 37°C. The reaction was stopped with 50µL/well of 0.37mM sulphuric acid. Absorbance was read at 492nm in an ELISA reader (Labsystems Multiskan MS 352, Finland).

Cell freezing. The hybrids were frozen according to Campbell et al. (1984), with modificatons. The freezing medium was HAT supplemented with 5% of DMSO. The hybridomas were kept at the Cell Bank of the Molecular and Cell Biology Laboratory, FUNED.

Cloning of the hybrid cells. The cloning of the hybridomas was performed by the limiting dilution procedure according to Harlow & Lane (1988) and the positive hybrids were frozen according to Campbell et al. (1984), with modificatons. The freezing medium was HAT supplemented with 5% of DMSO. The hybridomas were kept at the Cell Bank of the Molecular and Cell Biology Laboratory, FUNED.

Isotype characterization of monoclonal antibodies. The isotype characterization (classes and subclasses) of the produced MAbs were determined by ELISA as described (Harlow & Lane 1988). Briefly, after the addition of hybridoma supernatants to the plates (PoliSorp™, Nunc, Danmark), sensitized with sonicated antigen of *C. fetus* subsp. *venerealis* NCTC 10354 at a concentration of 3 ng/µL, 50µL of each specific MAbs against mouse IgA, IgM, IgG₁, IgG_{2a}, IgG_{2b} and IgG₃ (BD Biosciences Pharmingen, EUA) (2 µL/µL) were added per well and the plates were incubated at room temperature for 1h. The plates were then washed three times with PBS + 0.5% Tween 20, and 50µL of anti-rat IgG peroxidase conjugate (Sigma, USA), diluted 1:10,000 in incubation buffer, were added to each well, followed by incubation for 1 h at 37°C. Reactions were detected with OPD as previously described.

Characterization of the monoclonal antibody specificity. The specificity of the MAbs were assayed by indirect ELISA with sonicated antigens from *Arcobacter skirrowii* LMG 6621, *C. fetus* subsp. *fetus* ADRI 1812, *C. fetus* subsp. *venerealis* NCTC 10354, *C. lari* NCTC 11351, *C. sputorum* biovar sputorum LMG 6647. The MAbs reacting only against *C. fetus* antigens were also tested by western blot using sonicated antigens of *C. fetus* subsp. *venerealis* NCTC 10354 and *C. sputorum* biovar sputorum LMG 6647.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and Western blot. SDS-PAGE was performed as previously described (Laemmli 1970) using sonicated antigens from *C. fetus* subsp. *venerealis* NCTC 10354 and *C. sputorum* biovar sputorum LMG 6647. After electrophoresis, the separated antigens were transferred to a nitrocellulose membrane (Harlow & Lane 1988), at 100 V, for 1 h. Then, the nitrocellulose was washed three times with washing buffer (Tris saline plus 0.05% Tween 20 -TBST, pH 7.5), for 5 minutes at room temperature. The nitrocellulose membrane was blocked with TBS pH 7.5 + 1.0% Tween 20 for 1 h at room temperature. The nitrocellulose membrane was packed in Mini Protean®II Multi Screen (BioRad, EUA.) apparatus. The hybridoma supernatants (50µL) were added to the wells and incubated for 3 h at room temperature. After washing, the conjugate anti- mouse IgG peroxidase, at 1:5.000 diluiton in 0.1% TBST, was added and the membrane was incubated for 1 h at room temperature and developed with the substrate 3,3'-diaminobenzidine (DAB, Kit DAB SK-4100, Vector Laboratories, EUA) following the manufacturer's instructions.

RESULTS

Fusion with primed frozen spleen cells (Marusich 1988) resulted in no hybrids. The fusion with primed fresh spleen cells resulted in hybrids and the production of MAbs. Fourteen days after fusion, positive clones secreting antibodies anti-*Campylobacter fetus* subsp. *venerealis* were detected by indirect ELISA. After limiting dilution, 15 wells of the hybrid cells were positive for antibody against *C. fetus* subsp. *venerealis*.

Fourteen of the anti-*C. fetus* producing clones were of the IgG class, with subclasses IgG₁ (13 clones) and IgG_{2b} (1 clone), and one clone was of the IgM class (Table 1).

Results of the specificity characterization of the produced MAbs against antigens from *Arcobacter skirrowii* LMG 6621, *C. fetus* subsp. *fetus* serotype A ADRI 1812, *C. fetus* subsp. *venerealis* NCTC 10354, *C. lari* NCTC 11351 and *C. sputorum* biovar sputorum LMG 6647 are shown in Table 1.

The four producing clones of species-specific antibodies, M28 and M93, against *C. fetus* subsp. *venerealis*, and M7 and M94, specific for *C. fetus* subsp. *fetus*, were characterized by western blotting. The results showed that all clones recognized an immunogenic band with a molecular weight of approximately 148 kDa from the *C. fetus* subsp. *venerealis* antigen (Fig.1). No reactivity was observed with the antigen of *C. sputorum* biovar sputorum (data not shown).

DISCUSSION

The development in making of MAbs has greatly improved the production of specific and homogeneous antibodies. MAbs against microorganisms are important tools for the diagnosis and study of epidemiology and host-parasite relation in infectious diseases, such as those caused by *Campylobacter fetus* subsp. *venerealis*.

The protocol using primed frozen spleen cells (Marusich 1988) might constitute an interesting option for MAb production, as a sole spleen could be used at different times

Table 1. Specificity and isotype of monoclonal antibodies produced against *Campylobacter fetus* subsp. *venerealis* NCTC 10354

Clones	Strains ^a					Isotype
	LMG 6647	NCTC 10354	ADRI 1812	NCTC 11352	LMG 6621	
M2	- ^b	+ ^c	+	+	-	IgG ₁
M7	-	-	+	-	-	IgM
M21	-	+	+	+	-	IgG ₁
M22	-	+	+	+	-	IgG ₁
M28	-	+	-	-	-	IgG ₁
M37	-	+	+	+	-	IgG ₁
M39	-	+	+	+	-	IgG ₁
M41	-	+	+	+	-	IgG ₁
M42	-	+	+	+	-	IgG ₁
M44	-	+	-	+	+	IgG ₁
M45	-	+	+	+	+	IgG _{2b}
M60	-	+	+	+	-	IgG ₁
M76	-	+	+	+	-	IgG ₁
M93	-	+	-	-	-	IgG ₁
M94	-	-	+	-	-	IgG ₁

^a Strains: *Arcobacter skirrowii* LMG 6621, *C. fetus* subsp. *fetus* serotype A ADRI 1812, *C. fetus* subsp. *venerealis* NCTC 10354, *C. hyointestinalis* subsp. *hyointestinalis* LCDC 17398, *C. lari* NCTC11351, *C. sputorum* biovar sputorum LMG 6647; ^b "-" - no recognition; ^c "+" - recognition.




Fig.1. Western blotting for the characterization of monoclonal antibodies obtained from species-specific clones. Western blotting using sonicated *Campylobacter fetus* subsp. *venerealis* NCTC 10354 antigen (357 μ g per lane) and 2000 μ L of supernatant of culture clones M7 (lane 1), M28 (lane 2), M93 (lane 3), and M94 (lane 4). Reactions were performed using anti-mouse IgG peroxidase conjugate (Sigma, USA) and revealed by DAB kit (Vector Laboratories, USA). Arrow indicates the 148 kDa recognized band. Marker: Prestained Molecular Weight Standard mixture (Sigma, USA).

for fusion to obtain clones, therefore minimizing the use of mice. However, with the modified protocol used in the present study, no successful hybrid cell was obtained. Spleen cells were kept frozen for more than 2 years at -80°C in contrast with the prescribed storage at liquid nitrogen. Also, spleen cells were frozen in supplemented RPMI 1640 instead of HGD MEM supplemented medium. These differences could explain the failure in obtaining hybrid cells from frozen spleen cells. Many cell types stored at temperatures less than or around -80°C had their metabolism decreased, however to a level insufficient to halt all intracellular chemical reactions. This could result in the formation of intracellular ice crystals and damage of cellular structures, leading to cell death (Mazur 1984, Fowler & Toner 2005). However, for most of the cell types, freezing at -196°C halts cell metabolism as there is no liquid water and sufficient thermal energy for chemical reactions to occur, increasing the cell survival rates (Mazur 1984, Fowler & Toner 2005). RPMI 1640 medium has a lower concentration of glucose and sodium pyruvate that could represent readily available energy sources for unfrozen cells. Moreover, β -mercaptoethanol, insulin and the supernatant of P388D₁ cell culture (Bazin & Lemieur 1989) present in HGD MEM supplemented medium could have, respectively, facilitated cell fusion and stimulated energy metabolism and plasma cell proliferation. Irreversible damage to the cells could have occurred resulting in failure of hybrid cell production from frozen cells.

The fusion protocol used (Harlow & Lane 1988), performed with fresh spleen cells, resulted in antibody producing hybridomas, needing no feeder cells, due to their excellent growth and satisfactory number of cells to perform the cloning.

The predominance of MAb producing clones of the IgG class was expected due to the immunization scheme. The

use of complete and incomplete Freund adjuvants stimulate, respectively, predominant Th1 and Th2 responses with their high mainly associated production of IgG2a and IgG1 (Harlow & Lane 1988, Chang et al. 1998).

For the specificity studies, a panel of antigens was chosen based on the prevalences and habitats of bacteria from genera *Campylobacter* and *Arcobacter*. *C. sputorum* biovar *sputorum* is a commensal of the prepuce of bulls, being found in prepuce washings, which can mislead the diagnosis of *C. fetus* subsp. *venerealis* infection in bulls (Debruyne et al. 2008, Stoessel 1982). *C. fetus* subsp. *fetus* is the closest related to *C. fetus* subsp. *venerealis*, which inhabits the gastrointestinal tract of cattle and could also be found in the prepuce of bulls and in aborted fetuses from cattle (Blaser et al. 2008). The strain of *C. fetus* subsp. *fetus* used, ADRI 1812, is from serogroup A, to which all strains of *C. fetus* subsp. *venerealis* belong (Stoessel 1982). *C. lari* is an intestinal bacteria found in animals and humans. *Arcobacter skirrowii* can also be found in the prepuce of bulls being already isolated from aborted bovine fetus (Vandamme et al. 1992). Thus, results from the present study shows that MAbs obtained against *C. fetus* subsp. *venerealis* do not cross react with other bacteria of the *Campylobacteraceae* family that could be frequently present in the prepuce of bulls or in aborted bovine fetuses.

In the western blotting, no reactivity against the *C. sputorum* biovar *sputorum* was observed (data not shown) with the MAbs from the four species-specific clones, confirming again their specificity. Reactivity was observed against a protein of approximately 148 kDa from *C. fetus* subsp. *venerealis*. According to Dunn et al. (1987), proteins as the flagellum, with molecular mass of 63 kDa, proteins of the outer membrane of low molecular mass, varying from 43 to 44 kDa, and glycopolysaccharides, with 100 kDa, are important for immunogenicity against *C. fetus*. In *C. fetus*, immunologically relevant proteins are also found in a microcapsule-like structure named "S" layer that contains subunits of a protein known as SAP (surface array protein), formed by proteins of high molecular mass, varying from 97 to 149 kDa (Garcia et al. 1995). The "S" layer covers immunogenic structures of *C. fetus* including lipopolysaccharides and outer membrane proteins (Dunn et al. 1987, Garcia et al. 1995, Blaser et al. 2008). Changes in the "S" layer could explain the production of surface components with different antigenic specificities, which characterizes an evasion mechanism of the microorganism against the host immune system, allowing the *C. fetus* infection to persist longer (Blaser et al. 2008). Due to the SAP's high immunogenicity and the varying molecular weight, MAbs produced against *C. fetus* could probably be directed to SAP.

MAbs against *Campylobacter fetus* produced in the present study could constitute important tools for the development of diagnostic tests and also for characterization of *C. fetus* in studies of host-parasite relationship.

Acknowledgements. T.M. Alves and A.P. Lage were supported by fellowships from Conselho Nacional de Desenvolvimento Científico and Tecnológico (CNPq). This study received financial support from Fapemig and FEP-MVZ Coordenação Preventiva. All ADRI strains used in the present study were kindly supplied by Dr. B. Brooks. We thank C.P.T. de Brito and A. Fernandes for technical assistance.

REFERENCES

- Alves T.M., Stynen A.P.R., Miranda K.L. & Lage A.P. 2011. Campilobacteriose genital bovina e tricomonose genital bovina: epidemiologia, diagnóstico e controle. *Pesq. Vet. Bras.* 31(4):336-344.
- AVMA 2001. Report of the AVMA Panel on Euthanasia. *J. Am. Vet. Med. Assoc.* 218(5):669-696.
- Bazin R. & Lemieux R. 1989. Increased proportion of B cell hybridomas secreting monoclonal antibodies of desired specificity in cultures containing macrophage-derived hybridoma growth factor (IL-6). *J. Immunol. Meth.* 116:245-249.
- Blaser M.J., Newell D.G., Thompson S.A. & Zechner E.L. 2008. Pathogenesis of *Campylobacter fetus*, p.401-428. In: Nachamkin I., Szymanski C.M. & Blaser M.J. (Eds), *Campylobacter*. 3rd ed. ASM, Washington.
- Campbell A.M. 1984. *Monoclonal Antibody Technology*. Elsevier Science Publishers, Amsterdam. 265p.
- Chang J.C.C., Diveley J.P., Savary J.R. & Jensen F.C. 1998. Adjuvant activity of incomplete Freund's adjuvant. *Adv. Drug. Deliv. Rev.* 32:173-186.
- Coligan J.E., Kruisbeek A.M., Margulies D.H., Shevach E.M. & Strober W. 1998. *Current protocols in immunology*. Vol.1. John Wiley and Sons, New York.
- Debruyne L., Gevers D. & Vandamme P. 2008. Taxonomy of the family *Campylobacteraceae*, p.3-25. In: Nachamkin I., Szymanski C.M. & Blaser M.J. (Eds), *Campylobacter*. 3rd ed. ASM, Washington, DC.
- Dunn B.E., Blaser M.J. & Snyder E.L. 1987. Two-dimensional gel electrophoresis and immunoblotting of *Campylobacter* outer membrane proteins. *Infect. Immun.* 55(7):1564-1572.
- Fowler A. & Toner M. 2005. Cryo-injury and biopreservation. *Ann. N.Y. Acad. Sci.* 1066(1):119-135.
- Garcia M.M., Lutze-Wallace C.L., Denes A.S., Eaglesome M.D., Holst E. & Blaser M.J. 1995. Protein shift and antigenic variation in the S layer of *Campylobacter fetus* subsp. *venerealis* during bovine infection accompanied by genomic rearrangement of sapA homologs. *J. Bacteriol.* 177(8):1976-1980.
- Harlow E.D. & Lane D. 1988. *Antibody: A laboratory manual*. Cold Spring Harbor Laboratory, New York. 726p.
- Hurrell J.G.R. 1988. *Monoclonal Hybridoma Antibodies: Techniques and applications*. Cold Spring Harbor Laboratory, New York. 726p.
- Köhler G. & Milstein C. 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature.* 256:495-497.
- Laemmli U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-686.
- Lowry H.L., Rosebrough N.J. & Farr A.L. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193:265-275.
- Marusich M.F. 1988. Efficient hybridoma production using previously frozen splenocytes. *J. Immunol. Meth.* 144:155-159.
- Mazur P. 1984. Freezing of living cells: Mechanisms and implications. *Am. J. Physiol.* 247:125-142.
- Stoessel F. 1982. Las enfermedades venereas de los bovinos: trichomoniasis y vibriosis genital. *Acribia, Zaragoza.* 163p.
- Vandamme P., Vancanneyt M. & Pot B. 1992. Polyphasic taxonomy study of the emended genus *Arcobacter* with *Arcobacter butzleri* comb. nov. and *Arcobacter skirrowii* sp. nov., an aerotolerante bacterium isolated from veterinary specimens. *Int. J. Syst. Bacteriol.* 42:344-356.